

DESCRIPTIONS OF FOUR NEW SHELLS FROM THE PERSIAN GULF
AND BAY OF ZAILA.

By G. B. SOWERBY, F.L.S., etc.

Read June 8th, 1894.

PLATE XII. Figs. 20, 23, 25, 26.

1. CANCELLARIA PAUCICOSTATA, n.sp. Pl. XII. Fig. 26.

Testa angulata, utrinque conica, vix umbilicata, pallide fulva, obscure fusco fasciata; spira conica, gradata, medioeriter elevata; anfractus 6, convexi, embryonales lèves, sequentes obtuse angulati, longitudinaliter costati, liris numerosis confertis angustis alterne erassioribus eximie lamellatis spiraliter sculpti; anfractus ultimus costis paucis irregularibus instructus; apertura latiuscula, subtrigonalis, labrum leviter arcuaturn, fusco tintatum; columella rectiuscula, triplicata. Long. 17, lat. 11.50 mm.

Hab.—Persian Gulf.

An exquisitely sculptured shell, in which the angle of the body-whorl is lower than in the prevailing forms of the genus. The ribs on the body-whorl are few and irregularly placed. It would be difficult to assign the sectional or sub-generic position of this species, according to Dr. Jousseaume's arrangement. In form it seems intermediate between *Euelia*, H. and A. Adams, and *Scalptia*, Jousseaume; it has also some affinity with *Trigonostoma*, Blainville, as represented by *C. costata*, Gray, though it wants the umbilicus.

2. ANCILLA (MONOPTYGMA) UNIDENTATA, n.sp. Pl. XII. Fig. 20.

Testa oblongo subovata, crassa, polita, nitens, pallide fuscescens, hic illie castaneo strigata, postice castaneo latifasciata; spira exserta, obtusiusecula, callosa; apertura oblonga, postice angustiusecula, antice latior, trumeata; labrum lève, crassiuseculum; columella antice leviter contorta, plicata, postice callosa, plica unica dentiformi instructa. Long. 6, lat. 3 mm.

Hab.—Bay of Zaila.

A bright shining little shell, of which the spire is covered with a dark-brown enamel. It forms the second recent species of Lea's genus *Monoptygma*, characterized by a tooth-like plica above the middle of the columella.

3. NASSA ZAILENSIS, n.sp. Pl. XII. Fig. 25.

Testa acuminato-ovata, crassa, rugosa, griseo-albida, nigro-fusco fasciata; spira elata, acuta; anfractus 8, convexi, spiraliter subtilissime striati, costis numerosis nodulosis albo-articulatis instructi, nodulis obtusis, confertis; anfractus ultimus ventricosus, supra sulea latiuscula divisus; apertura subovata, intus costata, nigro-fusco fasciata, canali brevissimo; labrum politum, latiusculum, album, fusco pictum; columella brevis, alba, plicata, antice contorta. Long. 15, maj. lat. 8 mm.

Hab.—Bay of Zaila.

A prettily marked shell, in form somewhat like *N. vibex*, Say (sub-genus *Phrontis*, H. and A. Ad.).

4. *DONAX TOWNSENDI*, n.sp. Pl. XII. Fig. 23.

Testa sub-trigono oblonga, compressa, cuneata, valde inaequilateralis, polita, obscurissime radiatim striata, luteo-fusca, fasciis purpureis fuscisque irregulariter pieta; umbones acuti; margo dorsalis anticus rectiusculus, declivis, posticus truncatus, angulo postico obtusiusculo; area postangulari, bipartita, dimidio radiatim lirato, liris numerosis acutis hie illie elathrato, deinde laminis concentricis acutis laciniatis sculpta; pagina interna purpurea. Longit. 26, alt. 15·5 mm.

Var. a. Intus extusque lutea, postice pallide purpureo tineta.

Hab.—Persian Gulf.

This pretty species varies considerably in colour, the interior of the type being dark purple; some specimens are yellow within, and some nearly white, but these nearly all have a tinge of purple at the posterior end. The sculpture of the posterior area is rather peculiar, half the space behind the angle having sharp radiating liræ, and the other half concentric fringed laminæ, which, however, in most specimens more or less overlap the liræ, producing here and there a latticed sculpture.

EXPLANATION OF PLATE XII.

- Fig. 1. *Columbella (Amycla) planaxiformis.*
- „ 2. „ (*Mitrella* ?) *pupa.*
- „ 3. „ „ *semipieta.*
- „ 4. „ „ *venulata.*
- „ 5. „ (*Anachis*) *Sinensis.*
- „ 6. *Cingulina trisulcata.*
- „ 7. *Cerithiopsis multilirata.*
- „ 8. *Eulimella tenuis.*
- „ 9. *Turbanilla sinensis.*
- „ 10. „ *tenuistriata.*
- „ 11. *Odostomia subcarinata.*
- „ 12. *Alaba Hungerfordi.*
- „ 13. *Eulima producta.*
- „ 14. *Rissoia (Alvania) subcancellata.*
- „ 15. *Rissoina illustris.*
- „ 16. *Couthouyia acuta.*
- „ 17. *Adecorbis patula.*
- „ 18. *Minolia (Conotrochus) strigata.*
- „ 19. *Monilea rotundata.*
- „ 20. *Ancilla (Monoptygma) unidentata.*
- „ 21. *Psammobia simplex.*
- „ 22. *Tellina Hungerfordi.*
- „ 23. *Donax Townsendi.*
- „ 24. *Schizodentalium plurifissuratum*
- „ 25. *Nassa Zailensis.*
- „ 26. *Cancellaria paucicostata.*
- „ 27. *Plesiothyreus Newtoni.*